

ENCORE 603: A Phase II Randomized Study of Avelumab Plus Entinostat vs Avelumab Plus Placebo in Patients With Advanced and Recurrent Epithelial Ovarian Cancer

Karen Cadoo¹, Michael L. Meyers², Robert A. Burger³, Deborah Kay Armstrong⁴, Richard T. Penson⁵, Michael S. Gordon⁶, Gini F. Fleming⁷, John Moroney⁷, Erika Hamilton⁸, Linda R. Duska⁹, Robert Michael Wenham¹⁰, Serap Sankoh¹¹, Sue Brouwer¹¹, Peter Ordentlich¹¹, Carol Aghajanian¹, Ursula A. Matulonis¹²

¹Memorial Sloan Kettering Cancer Center, New York, NY; ²Syndax Pharmaceuticals, Inc., New York, NY; ³University of Pennsylvania, Philadelphia, PA;

⁴The Sidney Kimmel Comprehensive Cancer Center at Johns Hopkins University, Baltimore, MD; ⁵Harvard Medical School, Boston, MA;

⁶HonorHealth, Scottsdale, AZ; ⁷The University of Chicago Medicine, Chicago, IL; ⁸Sarah Cannon Research Institute, Nashville, TN; ⁹University of Virginia, Charlottesville, VA; ¹⁰Moffitt Cancer Center, Tampa, FL; ¹¹Syndax Pharmaceuticals, Inc., Waltham, MA; ¹²Dana-Farber Cancer Institute, Boston, MA

PRESENTED AT: **2019 ASCO**
ANNUAL MEETING

#ASCO19


Slides are the property of the author,
permission required for reuse.

PRESENTED BY: Karen Cadoo, MD

1

Background

- Avelumab is a humanized anti-PD-L1 IgG1 monoclonal antibody
- FDA approved in metastatic merkle cell carcinoma, advanced urothelial carcinoma and renal cancer



CD28, cluster of differentiation 28; MHC, major histocompatibility complex; PD-1, programmed cell death 1; PD-L1, programmed death ligand 1; PD-L2, programmed death ligand 2.
Adapted from: Postow MA, et al. *J Clin Oncol*. 2015;33(17):1974-1982.

Single-Agent Immune Checkpoint Inhibition Has Shown Limited Efficacy in Pretreated Ovarian Cancer

	KEYNOTE-028 ¹	JAVELIN Solid Tumor ²	KEYNOTE-100 ³	
Phase	1b	1b	2	
Patients, N	26	125	376	
Population	PD-L1–positive advanced	Recurrent/refractory	Advanced recurrent	
Prior lines of therapy, %	73% had ≥ 3 prior lines of therapy	65% had ≥ 3 prior lines of therapy	45% had ≥ 3 prior lines of therapy	
Treatment(s)	Pembrolizumab	Avelumab	Pembrolizumab	
ORR, %	11.5	9.6	8.0	
CR, %	3.8	0.8	1.9	
PR, %	7.7	8.8	6.1	
SD, %	26.9	42.4	29.3	
mPFS (95% CI), mo	1.9 (1.8–3.5)	2.6 (1.4–2.8)	2.1	
mOS (95% CI), mo	13.8 (6.7–18.8)	11.2 (8.7–15.4)	Cohort A [†] : NR (16.8–NR)	Cohort B [†] : 17.6 (13.3–NR)
Timing of scans	Baseline and every 8 weeks*	Baseline and every 6 weeks	Every 9 weeks [†]	


*After 6 months, scans were taken every 12 weeks. [†] Cohort A had 1-3 prior lines of therapy and Cohort B had 4-6 prior lines of therapy. [‡]Scans were taken every 12 weeks after week 54.

CI, confidence interval; CR, complete response; mOS, median overall survival; mPFS, median progression-free survival; NR, not reached; ORR, objective response rate; PD-L1, programmed death ligand-1; PR, partial response; SD, stable disease.

1. Varga A, et al. *Gynecol Oncol.* 2019;152(2):243-250. 2. Disis ML, et al. *JAMA Oncol.* 2019;5(3):393-401. 3. Matulonis UA, et al. *Ann Oncol.* 2019 May 2. doi: 10.1093/annonc/mdz135. [Epub ahead of print].

Potential for Immune Modulation with the Histone Deacetylase (HDAC) Inhibitor Entinostat


- Entinostat is an oral class I-selective histone deacetylase inhibitor
- Entinostat regulates the tumor microenvironment, immune inhibitory cells, induces a pro-inflammatory cascade, enhances antigen presentation
- Entinostat leads to additional beneficial effects on T_{eff} and NK cells



CTL, cytotoxic T lymphocytes; MDSC, myeloid-derived suppressor cells; NK, natural killer; TAM, tumor-associated macrophages; T_{eff}, effector T cell; Treg, regulatory T cell.
Orillion A, et al. *Clin Cancer Res.* 2017;23(17):5187-5201.

Preclinical Data has Demonstrated Synergy with Entinostat and anti PD-1 inhibition

- Entinostat leads to downregulation of immunosuppressive cell types (MDSC) in the tumor microenvironment




* $P < 0.05$. ** $P < 0.001$. [†] *In vivo* and *in vitro* studies were performed using Lewis Lung Carcinoma (LLC) cells.

Ab, antibody; Arg1, arginase 1; COX2, cytochrome oxidase subunit 2; iNOS, inducible nitric oxide synthase; MDSC, myeloid-derived suppressor cells. Orillion A, et al. *Clin Cancer Res.* 2017;23(17):5187-5201.

Preclinical Data has Demonstrated Synergy with Entinostat and anti PD-1 inhibition

- Entinostat leads to downregulation of immunosuppressive cell types (MDSC) in the tumor microenvironment






* $P < 0.05$. ** $P < 0.001$. [†] *In vivo* and *in vitro* studies were performed using Lewis Lung Carcinoma (LLC) cells.

Ab, antibody; Arg1, arginase 1; COX2, cytochrome oxidase subunit 2; iNOS, inducible nitric oxide synthase; MDSC, myeloid-derived suppressor cells. Orillion A, et al. *Clin Cancer Res.* 2017;23(17):5187-5201.

Preclinical Data has Demonstrated Synergy with Entinostat and anti PD-1 inhibition


- Entinostat leads to downregulation of immunosuppressive cell types (MDSC) in the tumor microenvironment
- Entinostat showed synergy with anti-PD-1 inhibition in preclinical models



* $P < 0.05$. ** $P < 0.001$. [†] *In vivo* and *in vitro* studies were performed using Lewis Lung Carcinoma (LLC) cells.


Ab, antibody; Arg1, arginase 1; COX2, cytochrome oxidase subunit 2; iNOS, inducible nitric oxide synthase; MDSC, myeloid-derived suppressor cells. Orillion A, et al. *Clin Cancer Res.* 2017;23(17):5187-5201.

ENCORE-603: Phase 2, Randomized, Double-Blind Study of Avelumab With or Without Entinostat in Patients With Advanced Epithelial Ovarian Cancer



Radiological disease assessments performed at baseline and every 6 weeks through week 36, thereafter every 8 weeks

ENCORE-603: Phase 2, Randomized, Double-Blind Study of Avelumab With or Without Entinostat in Patients With Advanced Epithelial Ovarian Cancer




Key eligibility:

- Patients must have progressive or recurrent ovarian cancer and 3-6 prior therapies
- Patients were agnostic to platinum sensitivity status
- No prior immunotherapy was allowed

Radiological disease assessments performed at baseline and every 6 weeks through week 36, thereafter every 8 weeks

ENCORE-603: Phase 2, Randomized, Double-Blind Study of Avelumab With or Without Entinostat in Patients With Advanced Epithelial Ovarian Cancer



Key eligibility:


- Patients must have progressive or recurrent ovarian cancer and 3-6 prior therapies
- Patients were agnostic to platinum sensitivity status
- No prior immunotherapy was allowed

Stratification:

- Presence of bulky disease (tumor $\geq 50\text{mm}$) vs not
- History of progression during primary platinum treatment (or within 1 month completion) vs not

Radiological disease assessments performed at baseline and every 6 weeks through week 36, thereafter every 8 weeks

ENCORE-603: Phase 2, Randomized, Double-Blind Study of Avelumab With or Without Entinostat in Patients With Advanced Epithelial Ovarian Cancer



Key eligibility:

- Patients must have progressive or recurrent ovarian cancer and 3-6 prior therapies
- Patients were agnostic to platinum sensitivity status
- No prior immunotherapy was allowed

Stratification:

- Presence of bulky disease (tumor $\geq 50\text{mm}$) vs not
- History of progression during primary platinum treatment (or within 1 month completion) vs not

Date of first patient in: July 11, 2017

Date of last patient in: April 17, 2018

Radiological disease assessments performed at baseline and every 6 weeks through week 36, thereafter every 8 weeks

Endpoints and Statistics

- Primary endpoint: Progression-Free Survival (PFS) based on RECIST 1.1
- Hypothesis: Avelumab + Entinostat would reduce the hazard of progression of disease or death by 43%, representing a 75% improvement in median PFS. 97 events (from 120 pts) provided 90% power with 1-sided significance level of 0.10
- Data cutoff date for final analysis of PFS: September 27, 2018
- Secondary endpoints: ORR by RECIST 1.1 and irRECIST, PFS by irRECIST, OS, CBR by RECIST 1.1 and irRECIST, AE, SAEs

Patient Demographics and Baseline Characteristics

	Avelumab + Placebo (N=41)	Avelumab + Entinostat (N=85)
	N (%)	N (%)
Median Age (range)	64 (45–79)	63 (43–82)
Race		
White	37 (90.2)	79 (92.9)
Other	4 (9.8)	6 (7.1)
ECOG Performance Status		
0	21 (51.2)	46 (54.1)
1	20 (48.8)	38 (44.7)
Body Mass Index		
< 30 kg/m ²	27 (65.9)	55 (64.7)
≥ 30 kg/m ²	12 (29.3)	24 (28.2)
Disease Stage at Initial Diagnosis		
II	4 (9.8)	10 (11.8)
III	25 (61.0)	45 (52.9)
IV	6 (14.6)	17 (20.0)

	Avelumab + Placebo (N=41)	Avelumab + Entinostat (N=85)
N (%)	N (%)	N (%)
Median Prior Therapies	4	4
Duration of Immediate Prior Therapy, Months		
Mean (SD)	3.5 (2.9)	4.1 (3.6)
Median (range)	2.8 (0.1–13.9)	3.5 (0.0–18.6)
Presence of Bulky Disease (>50mm)		
Yes	11 (26.8)	25 (29.4)
No	30 (73.2)	60 (70.6)
Histologies		
Serous	36 (87.8)	69 (81.2)
Endometroid	1 (2.4)	4 (4.7)
Clear cell	1 (2.4)	3 (3.5)
Other	0 (0.0)	5 (5.9)

ECOG, Eastern Cooperative Oncology Group.

Patient Demographics and Baseline Characteristics

	Avelumab + Placebo (N=41)	Avelumab + Entinostat (N=85)
	N (%)	N (%)
Median Age (range)	64 (45–79)	63 (43–82)
Race		
White	37 (90.2)	79 (92.9)
Other	4 (9.8)	6 (7.1)
ECOG Performance Status		
0	21 (51.2)	46 (54.1)
1	20 (48.8)	38 (44.7)
Body Mass Index		
< 30 kg/m ²	27 (65.9)	55 (64.7)
≥ 30 kg/m ²	12 (29.3)	24 (28.2)
Disease Stage at Initial Diagnosis		
II	4 (9.8)	10 (11.8)
III	25 (61.0)	45 (52.9)
IV	6 (14.6)	17 (20.0)

	Avelumab + Placebo (N=41)	Avelumab + Entinostat (N=85)
N (%)	N (%)	N (%)
Median Prior Therapies	4	4
Duration of Immediate Prior Therapy, Months		
Mean (SD)	3.5 (2.9)	4.1 (3.6)
Median (range)	2.8 (0.1–13.9)	3.5 (0.0–18.6)
Presence of Bulky Disease (>50mm)		
Yes	11 (26.8)	25 (29.4)
No	30 (73.2)	60 (70.6)
Histologies		
Serous	36 (87.8)	69 (81.2)
Endometroid	1 (2.4)	4 (4.7)
Clear cell	1 (2.4)	3 (3.5)
Other	0 (0.0)	5 (5.9)

ECOG, Eastern Cooperative Oncology Group.

Patient Demographics and Baseline Characteristics

	Avelumab + Placebo (N=41)	Avelumab + Entinostat (N=85)
	N (%)	N (%)
Median Age (range)	64 (45–79)	63 (43–82)
Race		
White	37 (90.2)	79 (92.9)
Other	4 (9.8)	6 (7.1)
ECOG Performance Status		
0	21 (51.2)	46 (54.1)
1	20 (48.8)	38 (44.7)
Body Mass Index		
< 30 kg/m ²	27 (65.9)	55 (64.7)
≥ 30 kg/m ²	12 (29.3)	24 (28.2)
Disease Stage at Initial Diagnosis		
II	4 (9.8)	10 (11.8)
III	25 (61.0)	45 (52.9)
IV	6 (14.6)	17 (20.0)

	Avelumab + Placebo (N=41)	Avelumab + Entinostat (N=85)
N (%)	N (%)	N (%)
Median Prior Therapies	4	4
Duration of Immediate Prior Therapy, Months		
Mean (SD)	3.5 (2.9)	4.1 (3.6)
Median (range)	2.8 (0.1–13.9)	3.5 (0.0–18.6)
Presence of Bulky Disease (>50mm)		
Yes	11 (26.8)	25 (29.4)
No	30 (73.2)	60 (70.6)
Histologies		
Serous	36 (87.8)	69 (81.2)
Endometroid	1 (2.4)	4 (4.7)
Clear cell	1 (2.4)	3 (3.5)
Other	0 (0.0)	5 (5.9)

ECOG, Eastern Cooperative Oncology Group.

Patient Demographics and Baseline Characteristics

	Avelumab + Placebo (N=41)	Avelumab + Entinostat (N=85)
	N (%)	N (%)
Median Age (range)	64 (45–79)	63 (43–82)
Race		
White	37 (90.2)	79 (92.9)
Other	4 (9.8)	6 (7.1)
ECOG Performance Status		
0	21 (51.2)	46 (54.1)
1	20 (48.8)	38 (44.7)
Body Mass Index		
< 30 kg/m ²	27 (65.9)	55 (64.7)
≥ 30 kg/m ²	12 (29.3)	24 (28.2)
Disease Stage at Initial Diagnosis		
II	4 (9.8)	10 (11.8)
III	25 (61.0)	45 (52.9)
IV	6 (14.6)	17 (20.0)

	Avelumab + Placebo (N=41)	Avelumab + Entinostat (N=85)
N (%)	N (%)	N (%)
Median Prior Therapies	4	4
Duration of Immediate Prior Therapy, Months		
Mean (SD)	3.5 (2.9)	4.1 (3.6)
Median (range)	2.8 (0.1–13.9)	3.5 (0.0–18.6)
Presence of Bulky Disease (>50mm)		
Yes	11 (26.8)	25 (29.4)
No	30 (73.2)	60 (70.6)
Histologies		
Serous	36 (87.8)	69 (81.2)
Endometrioid	1 (2.4)	4 (4.7)
Clear cell	1 (2.4)	3 (3.5)
Other	0 (0.0)	5 (5.9)

ECOG, Eastern Cooperative Oncology Group.

Patient Demographics and Baseline Characteristics

	Avelumab + Placebo (N=41)	Avelumab + Entinostat (N=85)
	N (%)	N (%)
Median Age (range)	64 (45–79)	63 (43–82)
Race		
White	37 (90.2)	79 (92.9)
Other	4 (9.8)	6 (7.1)
ECOG Performance Status		
0	21 (51.2)	46 (54.1)
1	20 (48.8)	38 (44.7)
Body Mass Index		
< 30 kg/m ²	27 (65.9)	55 (64.7)
≥ 30 kg/m ²	12 (29.3)	24 (28.2)
Disease Stage at Initial Diagnosis		
II	4 (9.8)	10 (11.8)
III	25 (61.0)	45 (52.9)
IV	6 (14.6)	17 (20.0)

	Avelumab + Placebo (N=41)	Avelumab + Entinostat (N=85)
N (%)	N (%)	N (%)
Median Prior Therapies	4	4
Duration of Immediate Prior Therapy, Months		
Mean (SD)	3.5 (2.9)	4.1 (3.6)
Median (range)	2.8 (0.1–13.9)	3.5 (0.0–18.6)
Presence of Bulky Disease (>50mm)		
Yes	11 (26.8)	25 (29.4)
No	30 (73.2)	60 (70.6)
Histologies		
Serous	36 (87.8)	69 (81.2)
Endometroid	1 (2.4)	4 (4.7)
Clear cell	1 (2.4)	3 (3.5)
Other	0 (0.0)	5 (5.9)

ECOG, Eastern Cooperative Oncology Group.

Patient Demographics and Baseline Characteristics

	Avelumab + Placebo (N=41)	Avelumab + Entinostat (N=85)
	N (%)	N (%)
Median Age (range)	64 (45–79)	63 (43–82)
Race		
White	37 (90.2)	79 (92.9)
Other	4 (9.8)	6 (7.1)
ECOG Performance Status		
0	21 (51.2)	46 (54.1)
1	20 (48.8)	38 (44.7)
Body Mass Index		
< 30 kg/m ²	27 (65.9)	55 (64.7)
≥ 30 kg/m ²	12 (29.3)	24 (28.2)
Disease Stage at Initial Diagnosis		
II	4 (9.8)	10 (11.8)
III	25 (61.0)	45 (52.9)
IV	6 (14.6)	17 (20.0)

ECOG, Eastern Cooperative Oncology Group.

	Avelumab + Placebo (N=41)	Avelumab + Entinostat (N=85)
N (%)	N (%)	N (%)
Median Prior Therapies	4	4
Duration of Immediate Prior Therapy, Months		
Mean (SD)	3.5 (2.9)	4.1 (3.6)
Median (range)	2.8 (0.1–13.9)	3.5 (0.0–18.6)
Presence of Bulky Disease (>50mm)		
Yes	11 (26.8)	25 (29.4)
No	30 (73.2)	60 (70.6)
Histologies		
Serous	36 (87.8)	69 (81.2)
Endometrioid	1 (2.4)	4 (4.7)
Clear cell	1 (2.4)	3 (3.5)
Other	0 (0.0)	5 (5.9)

Patient Status

N (%)	Avelumab + Placebo (N=41)	Avelumab + Entinostat (N=85)
Patients Randomized	41 (100)	85 (100)
Safety Population	40 (97.6)	85 (100)
Treatment Status		
On study treatment	4 (10.0)	9 (10.6)
Discontinued study therapy	36 (90.0)	76 (89.4)
Progressive disease (RECIST 1.1)	21 (52.5)	46 (54.1)
Progressive disease (irRECIST)	2 (5.0)	4 (4.7)
Clinical progression	1 (2.5)	5 (5.9)
Adverse event	7 (17.5)	17 (20.0)
Withdrawal of consent	3 (7.5)	1 (1.2)
Other	2 (5.0)	3 (3.5)
Stayed on treatment after RECIST 1.1 PD	10 (24.4)	18 (21.2)

Patient Status

N (%)	Avelumab + Placebo (N=41)	Avelumab + Entinostat (N=85)
Patients Randomized	41 (100)	85 (100)
Safety Population	40 (97.6)	85 (100)
Treatment Status		
On study treatment	4 (10.0)	9 (10.6)
Discontinued study therapy	36 (90.0)	76 (89.4)
Progressive disease (RECIST 1.1)	21 (52.5)	46 (54.1)
Progressive disease (irRECIST)	2 (5.0)	4 (4.7)
Clinical progression	1 (2.5)	5 (5.9)
Adverse event	7 (17.5)	17 (20.0)
Withdrawal of consent	3 (7.5)	1 (1.2)
Other	2 (5.0)	3 (3.5)
Stayed on treatment after RECIST 1.1 PD	10 (24.4)	18 (21.2)

Patient Status

N (%)	Avelumab + Placebo (N=41)	Avelumab + Entinostat (N=85)
Patients Randomized	41 (100)	85 (100)
Safety Population	40 (97.6)	85 (100)
Treatment Status		
On study treatment	4 (10.0)	9 (10.6)
Discontinued study therapy	36 (90.0)	76 (89.4)
Progressive disease (RECIST 1.1)	21 (52.5)	46 (54.1)
Progressive disease (irRECIST)	2 (5.0)	4 (4.7)
Clinical progression	1 (2.5)	5 (5.9)
Adverse event	7 (17.5)	17 (20.0)
Withdrawal of consent	3 (7.5)	1 (1.2)
Other	2 (5.0)	3 (3.5)
Stayed on treatment after RECIST 1.1 PD	10 (24.4)	18 (21.2)


Patient Status

N (%)	Avelumab + Placebo (N=41)	Avelumab + Entinostat (N=85)
Patients Randomized	41 (100)	85 (100)
Safety Population	40 (97.6)	85 (100)
Treatment Status		
On study treatment	4 (10.0)	9 (10.6)
Discontinued study therapy	36 (90.0)	76 (89.4)
Progressive disease (RECIST 1.1)	21 (52.5)	46 (54.1)
Progressive disease (irRECIST)	2 (5.0)	4 (4.7)
Clinical progression	1 (2.5)	5 (5.9)
Adverse event	7 (17.5)	17 (20.0)
Withdrawal of consent	3 (7.5)	1 (1.2)
Other	2 (5.0)	3 (3.5)
Stayed on treatment after RECIST 1.1 PD	10 (24.4)	18 (21.2)

Patient Status


N (%)	Avelumab + Placebo (N=41)	Avelumab + Entinostat (N=85)
Patients Randomized	41 (100)	85 (100)
Safety Population	40 (97.6)	85 (100)
Treatment Status		
On study treatment	4 (10.0)	9 (10.6)
Discontinued study therapy	36 (90.0)	76 (89.4)
Progressive disease (RECIST 1.1)	21 (52.5)	46 (54.1)
Progressive disease (irRECIST)	2 (5.0)	4 (4.7)
Clinical progression	1 (2.5)	5 (5.9)
Adverse event	7 (17.5)	17 (20.0)
Withdrawal of consent	3 (7.5)	1 (1.2)
Other	2 (5.0)	3 (3.5)
Stayed on treatment after RECIST 1.1 PD	10 (24.4)	18 (21.2)

Primary Endpoint Progression-Free Survival RECIST 1.1



Radiological disease assessments performed at baseline and every 6 weeks through week 36, and after week 36 every 8 weeks

Subgroup Analysis




Secondary Endpoints: Objective Response Rate and Clinical Benefit Rate


Endpoint	Avelumab + Placebo (N=41)	Avelumab + Entinostat (N=85)
	N (%)	N (%)
Objective Response Rate (RECIST 1.1)	2 (4.9)	5 (5.9)
Exact 95% confidence interval	(0.6, 16.5)	(1.9, 13.2)
Clinical Benefit Rate (CR+PR+SD $\geq$ 6 months)	4 (9.8)	10 (11.8)
Exact 95% confidence interval	(2.7, 23.1)	(5.8, 20.6)
Stable Disease $\geq$ 4 months	3 (7.3)	15 (17.6)
Exact 95% confidence interval	(1.5, 19.9)	(10.2, 27.4)

*Per protocol, clinical benefit rate was defined as complete response + partial response + stable disease $\geq$ 6 months.


Select Patients With Durable Benefit




Select Patients With Durable Benefit



Select Patients With Durable Benefit



Select Patients With Durable Benefit



Safety Summary

Patients with:	Avelumab + Placebo (N=40)	Avelumab + Entinostat (N=85)
	N (%)	N (%)
Any related TEAE	31 (77.5)	79 (92.9)
Related grade 3/4 TEAE	4 (10.0)	35 (41.2)
TEAE that led to any study drug being permanently discontinued	7 (17.5)	18 (21.2)
TEAE that led to any study treatment being dose modified	11 (27.5)	50 (58.8)
Deaths on study, all unrelated	4 (10.0)	6 (7.0)

TEAE- treatment emergent adverse event

Safety Summary

Patients with:	Avelumab + Placebo (N=40)	Avelumab + Entinostat (N=85)
	N (%)	N (%)
Any related TEAE	31 (77.5)	79 (92.9)
Related grade 3/4 TEAE	4 (10.0)	35 (41.2)
TEAE that led to any study drug being permanently discontinued	7 (17.5)	18 (21.2)
TEAE that led to any study treatment being dose modified	11 (27.5)	50 (58.8)
Deaths on study, all unrelated	4 (10.0)	6 (7.0)

TEAE- treatment emergent adverse event



Grade 3/4 Adverse Events

Grade 3/4 Related Adverse Events in > 1 Patients in Either Treatment Arm	Avelumab + Placebo (N=40)	Avelumab + Entinostat (N=85)
	N (%)	N (%)
Patients With At Least One Treatment Emergent Adverse Event	4 (10.0)	35 (41.2)
Neutropenia	0	11 (12.9)
Fatigue	1 (2.5)	8 (9.4)
Anemia	0	4 (4.7)
Hyponatremia	0	4 (4.7)
Nausea	1 (2.5)	2 (2.4)
Hypocalcemia	0	2 (2.4)
Thrombocytopenia	0	2 (2.4)
Dehydration	1 (2.5)	1 (1.2)
Infusion related reaction	1 (2.5)	1 (1.2)
Vomiting	1 (2.5)	1 (1.2)

Immune-Related Adverse Events

Immune Related Adverse Events > 1 Patient All Grade	Avelumab + Placebo (N=40)		Avelumab + Entinostat (N=85)	
	All Grades	Grade 3/4	All Grades	Grade 3/4
	N (%)	N (%)	N (%)	N (%)
Hypothyroidism	3 (7.5)	0	3 (3.5)	0
Pruritus	1 (2.5)	0	3 (3.5)	0
Colitis	0	0	3 (3.5)	1 (1.2)
Pneumonitis	0	0	3 (3.5)	0
Rash	1 (2.5)	0	2 (2.4)	0
Rash-maculo-papular	0	0	2 (2.4)	0

Entinostat Plus Avelumab Reduced Circulating MDSC



- Entinostat plus Avelumab resulted in a statistically significant decrease in MDSC at C2D8 and C4D1 relative to C1D1-Predose, $p < 0.01$
- No significant changes of MDSC levels in Avelumab monotherapy treatment arm

MDSC, myeloid-derived suppressor cells; ns, not significant.

Conclusions

- Despite good rationale and demonstrating suppression of myeloid derived suppressor cells, the addition of Entinostat did not improve outcomes over Avelumab alone for patients with heavily pretreated advanced ovarian cancer
- As with other immunotherapy studies in ovarian cancer, there are clearly patients who benefit
- Overall survival data remains immature and correlative studies are ongoing, we are eagerly awaiting further data which may help us to understand which patients with ovarian cancer benefit from immunotherapy

Conclusions

- There was no new safety signal and the addition of Entinostat did not lead to an excess of immune-related adverse events
- Only approximately 20% of patients remained on therapy beyond RECIST 1.1 progression of disease
- This remains an area of huge unmet need, we enrolled 126 advanced ovarian cancer patients in 9 months

Acknowledgments

- We thank the patients and their families
- Co-Investigators and study staff

This study was sponsored by Syndax Pharmaceuticals, Inc., in collaboration with Merck KGaA, Darmstadt, Germany and Pfizer. Editorial assistance was provided by Kristen Evaul PhD. This assistance was funded by Syndax Pharmaceuticals, Inc.

PRESENTED AT: **2019 ASCO**
ANNUAL MEETING

#ASCO19
Slides are the property of the author,
permission required for reuse.

PRESENTED BY: Karen Cadoo, MD

38